

# Nayzilam®<sup>IV</sup> (midazolam) nasal spray



## UNDERSTANDING NAYZILAM: A READY-TO-USE TREATMENT FOR SEIZURE CLUSTERS

NAYZILAM is a prescription medicine used for the short-term treatment of seizure clusters (also known as “acute repetitive seizures”) in patients 12 years of age and older.

### IMPORTANT SAFETY INFORMATION

NAYZILAM is a benzodiazepine medicine. Taking benzodiazepines with opioid medicines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, breathing problems (respiratory depression), coma, and death. The most common adverse reactions ( $\geq 5\%$  in any NAYZILAM treatment group) were somnolence, headache, nasal discomfort, throat irritation, and rhinorrhea.

# CONTENTS

## GETTING STARTED WITH NAYZILAM

PAGE 3

## USING NAYZILAM

PAGE 8

## UNDERSTANDING SEIZURE CLUSTERS

PAGE 4

## HOW TO GIVE NAYZILAM

PAGE 9

## WHAT IS NAYZILAM?

PAGE 6

## PATIENT SUPPORT

PAGE 14

## HOW NAYZILAM CAN HELP

PAGE 7

## SAVINGS PROGRAM

PAGE 16



Please see additional Important Safety Information on pages 10-13 and the enclosed Patient Medication Guide in this brochure's pocket. Visit [www.NAYZILAM.com](http://www.NAYZILAM.com) to learn more about treatment with NAYZILAM.

# GETTING STARTED WITH NAYZILAM

Your healthcare provider has prescribed NAYZILAM® (midazolam) nasal spray, CIV as a way to help you or a loved one manage seizure clusters when they happen. This guide will help you to understand:

- What seizure clusters are
- How to manage them
- How NAYZILAM can help treat them quickly



**PLEASE TALK TO YOUR HEALTHCARE PROVIDER IF YOU HAVE ANY QUESTIONS OR CONCERNS ABOUT NAYZILAM.**

Living with seizure clusters can be worrisome, but now you can have the peace of mind that comes from having a fast-acting rescue medication in a ready-to-use nasal spray.

*I would like to see a treatment that would reduce the frequency of seizure clusters in a day. This would open up for my brother many opportunities he currently does not have, such as being able to work, and greatly improve his outlook on life and the future.*

- Caregiver

## IMPORTANT SAFETY INFORMATION

NAYZILAM may cause serious side effects, including: interaction with other central nervous system depressants, serious breathing problems and excessive sleepiness, suicidal thoughts or actions, impaired mental alertness including memory problems, and increase in eye pressure in people with acute narrow-angle glaucoma. The most common side effects of NAYZILAM include sleepiness, nasal discomfort, headache, throat irritation, and runny nose.

# UNDERSTANDING SEIZURE CLUSTERS

Seizure clusters are disruptive events that can leave you feeling worried, frustrated, or helpless.



**IF YOU EXPERIENCE SEIZURE CLUSTERS,  
YOU ARE NOT ALONE.**

## WHAT ARE SEIZURE CLUSTERS?

Seizure clusters are periods of increased seizure activity, which is having two or more seizures in a 24-hour period.

You may experience seizures, but *seizure clusters* are different from your usual seizure pattern. This can mean having several seizures in a row or they can be hours apart. Your healthcare provider can help you to identify seizure clusters and when and how to use a rescue medication to treat them.

You may have heard of seizure clusters, or your healthcare provider may call them something else, such as:

- Acute repetitive seizures
- Serial seizures
- Recurrent seizures
- Multiple seizures
- Crescendo seizures
- Seizure flurries

**SEIZURE CLUSTERS = 2 OR MORE SEIZURES IN 24 HOURS**

## WHY IT'S URGENT TO TREAT SEIZURE CLUSTERS

Seizure clusters may happen without warning and can lead to serious health consequences, such as:

- Injuries
- Hospitalization
- Longer seizures or more dangerous activity, called status epilepticus, which is a seizure lasting longer than five minutes

That's why it's important to have a rescue medication on hand during a seizure cluster emergency, and to use it as soon as possible.

What matters most is having the reassurance of a rescue treatment that:

- Works fast to stop a seizure cluster
- Allows you to feel like yourself again
- Prevents additional seizures within the cluster
- Can be given by anyone, anywhere

A rescue medication is different from the daily epilepsy medications you already take to manage your epilepsy. Keep taking your daily seizure medication as directed by your healthcare provider, and use your prescribed rescue medication when you need it to treat a seizure cluster.

**DON'T LET ONE SEIZURE  
LEAD TO ANOTHER.**

**TREAT SEIZURE CLUSTERS  
QUICKLY SO THEY DON'T TURN  
INTO A DANGEROUS THREAT.**

## WHAT IS NAYZILAM?

NAYZILAM® (midazolam) nasal spray, CIV is a prescription medicine used for the short-term treatment of seizure clusters (also known as acute repetitive seizures) in patients 12 years of age and older. It is a rescue medication in the form of a nasal spray.

Do not use NAYZILAM if you have an eye problem called acute narrow-angle glaucoma.

You don't have to wait to treat a seizure cluster with NAYZILAM. This nasal spray can be given during or immediately after the first seizure if a cluster occurs, even if the person is unconscious—no inhaling is necessary.

Prior to using NAYZILAM, the person giving it should know how to identify seizure clusters and how to use this rescue medication appropriately.

NAYZILAM is different from daily epilepsy medications prescribed to manage epilepsy. Keep taking your daily seizure medication as directed by your healthcare provider, and use NAYZILAM when you need to treat a seizure cluster.

To learn more about NAYZILAM, please visit [www.NAYZILAM.com](http://www.NAYZILAM.com).



### IMPORTANT SAFETY INFORMATION

NAYZILAM is a benzodiazepine medicine. Taking benzodiazepines with opioid medicines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, breathing problems (respiratory depression), coma, and death. NAYZILAM may cause serious breathing problems and excessive sleepiness (sedation).

**Nayzilam**®  
(midazolam) nasal spray

Please see additional Important Safety Information on pages 10-13 and the enclosed Patient Medication Guide in this brochure's pocket. Visit [www.NAYZILAM.com](http://www.NAYZILAM.com) to learn more about treatment with NAYZILAM.

## HOW NAYZILAM CAN HELP

Now you have the reassurance of NAYZILAM within reach to help manage seizure clusters quickly and get back to your day.

**IN A STUDY, NAYZILAM HELPED THE MAJORITY OF PATIENTS TO STOP A SEIZURE CLUSTER FAST AND RETURN TO FUNCTION WITHIN 90 MINUTES.**

NAYZILAM may cause serious side effects, including impaired mental alertness including memory problems.

The most common side effects reported with NAYZILAM use include:

- Sleepiness
- Headache
- Nasal discomfort
- Throat irritation
- Runny nose

These are not all the possible side effects of NAYZILAM. Call your healthcare provider for medical advice about side effects.



**STOP SEIZURE CLUSTERS SO THEY DON'T STOP YOU. WITH NAYZILAM WITHIN REACH, NOW YOU CAN BE READY TO TREAT THEM QUICKLY.**

NAYZILAM MAY HELP YOU TO:

**STOP SEIZURE CLUSTERS WITHIN MINUTES**

Prevent further seizure activity for up to 6 hours

**GET BACK TO FEELING LIKE YOURSELF WITHIN HOURS**

**KEEP MORE SEIZURES FROM RETURNING FOR 24 HOURS**

## USING NAYZILAM

 EACH BOX CONTAINS TWO SINGLE-USE NASAL SPRAY DEVICES

 FIRST DOSE IS ONE 5 MG NASAL SPRAY (ONLY ONE DOSE MAY BE NEEDED)

 STORE AT ROOM TEMPERATURE



### ABOUT USING NAYZILAM

It's important to know a few things before you or someone else administers NAYZILAM® (midazolam) nasal spray, CIV, so everyone can feel more comfortable using it.

- NAYZILAM is not intended to be taken daily. Use exactly as your healthcare provider tells you to use it (in the nose, never more than two doses). If you use too much of this medication, call your healthcare provider or go to the nearest emergency room right away.
- If the seizure cluster is continuing 10 minutes after the first dose, a second dose may be used if you have been told to do so by your healthcare provider. If a second dose is used, give the second dose in the other nostril.
- If the seizures do not stop after the nasal spray is used, get emergency medical help right away.
- Do not use this medication for more than one seizure cluster episode every 3 days and no more than five seizure cluster episodes per month.

Please see the enclosed Medication Guide for more information.

 PLEASE TALK TO YOUR HEALTHCARE PROVIDER IF YOU HAVE ANY QUESTIONS OR CONCERNS ABOUT NAYZILAM.

Nayzilam®  
(midazolam) nasal spray

Please see additional Important Safety Information on pages 10-13 and the enclosed Patient Medication Guide in this brochure's pocket. Visit [www.NAYZILAM.com](http://www.NAYZILAM.com) to learn more about treatment with NAYZILAM.

## HOW TO GIVE NAYZILAM

NAYZILAM is a ready-to-use nasal spray that can be given by anyone, whenever seizure clusters occur.

 AFTER PEELING OPEN THE BLISTER PACK, JUST FOLLOW THESE STEPS.

### HOLD

Hold the nasal spray device with your thumb on the plunger and your middle and index fingers on each side of the nozzle.



DO NOT PRESS  
THE PLUNGER  
YET.

### PLACE


Place the tip of the nozzle into one nostril until your fingers are against the bottom of the patient's nose.



### PRESS

Press the plunger firmly.



 REMEMBER TO THROW AWAY (DISPOSE OF) THE NASAL SPRAY UNIT AND BLISTER PACKAGING.

These are not the complete instructions for use. Please read the enclosed full Instructions for Use in this brochure's pocket.

**NOW YOU CAN FEEL READY TO TREAT SEIZURE CLUSTERS ANYTIME WITH NAYZILAM.**



# IMPORTANT SAFETY INFORMATION

Important Safety Information you should know about NAYZILAM® (midazolam) nasal spray, CIV

What is the most important information I should know about NAYZILAM?

NAYZILAM is a benzodiazepine medicine. Taking benzodiazepines with opioid medicines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, breathing problems (respiratory depression), coma, and death.

NAYZILAM may cause serious breathing problems and excessive sleepiness (sedation). Get emergency medical help right away if any of the following occur:

- unusually shallow or slowed breathing
- stop breathing, which may lead to your heart stopping
- unusually excessive sleepiness

Like other antiepileptic drugs, NAYZILAM may cause suicidal thoughts or actions in a very small number of people, about 1 in 500. Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- feeling agitated or restless
- acting aggressive, being angry, or violent
- attempts to commit suicide
- panic attacks
- acting on dangerous impulses
- trouble sleeping (insomnia)
- an extreme increase in activity and talking (mania)
- new or worse anxiety
- new or worse irritability
- other unusual changes in behavior or mood



How can I watch for early symptoms of suicidal thoughts or actions?

- Pay attention to any changes, especially sudden changes in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms. Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

What is NAYZILAM?

- NAYZILAM is a prescription medicine used for the short-term treatment of seizure clusters (also known as “acute repetitive seizures”) in patients 12 years of age and older.
- **NAYZILAM is a federal controlled substance (CIV) because it can be abused or lead to dependence.** Keep NAYZILAM in a safe place to prevent misuse and abuse. Selling or giving away NAYZILAM may harm others and is against the law. Tell your healthcare provider if you have abused or been dependent on alcohol, prescription drugs, or street drugs.
- It is not known if NAYZILAM is safe and effective in children under 12 years of age.

Do not use NAYZILAM if you:

- are allergic to midazolam.
- have an eye problem called acute narrow-angle glaucoma.

Before you use NAYZILAM, tell your healthcare provider about all your medical conditions, including if you:

- have a history of depression, mood problems or suicidal thoughts or behavior.
- have asthma, emphysema, bronchitis, chronic obstructive pulmonary disease, or other breathing problems.
- have kidney or liver problems.
- have congestive heart failure.
- have a history of drug or alcohol abuse.
- are pregnant or plan to become pregnant. NAYZILAM may harm your unborn baby.
- are breastfeeding or plan to breastfeed.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

#### How should I use NAYZILAM® (midazolam) nasal spray, CIV?

- Use NAYZILAM in the nose only.
- **Use NAYZILAM exactly as your healthcare provider tells you to use it** and follow the Instructions for Use.
- Your healthcare provider has prescribed NAYZILAM to treat a type of seizure called a “seizure cluster”.
- If the seizure cluster is continuing **10 minutes** after the first dose of NAYZILAM, a second dose of NAYZILAM **may be used if you have been told to do so by your healthcare provider**.
- If a second dose of NAYZILAM is used, give the second dose in the other nostril.
- **Do not** give more than 2 doses of NAYZILAM to treat a seizure cluster.
- **If the seizures do not stop** after NAYZILAM is used, get emergency medical help right away.
- **Do not** use NAYZILAM for more than one seizure cluster episode every 3 days.  
**Do not** use NAYZILAM for more than five seizure cluster episodes per month.
- If benzodiazepines are stopped after a person takes them daily, they can cause withdrawal symptoms. Stopping benzodiazepines suddenly can cause seizures that will not stop (status epilepticus), hearing or seeing things that are not there (hallucinations), shaking, nervousness, and stomach and muscle cramps. NAYZILAM is not intended to be taken daily.
- If you use too much NAYZILAM, call your healthcare provider or go to the nearest emergency room right away.

#### What should I avoid while using NAYZILAM?

- **Do not** drive, operate machinery, or do other activities that require mental alertness until you know how NAYZILAM affects you.
- **Do not** drink alcohol or take opioid medicines or other medicines that make you sleepy or dizzy while taking NAYZILAM until you talk to your healthcare provider.



#### What are the possible side effects of NAYZILAM?

NAYZILAM may cause serious side effects, including:

- See “**What is the most important information I should know about NAYZILAM?**” for information on serious side effects of:
  - interaction with other central nervous system depressants
  - serious breathing problems and excessive sleepiness
  - suicidal thoughts or actions
- **Impaired mental alertness including memory problems.**  
See “**What should I avoid while using NAYZILAM?**”
- **Increase in eye pressure in people with acute narrow-angle glaucoma.**  
See “**Do not take NAYZILAM if you:**”

The most common side effects of NAYZILAM include:

- sleepiness
- nasal discomfort
- headache
- throat irritation
- runny nose

These are not all the possible side effects of NAYZILAM. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see additional patient information in the Medication Guide. This information does not take the place of talking with your healthcare provider about your condition or your treatment. For more information, go to [www.NAYZILAM.com](http://www.NAYZILAM.com) or call 1-844-599-2273.

# PATIENT SUPPORT

## GET A TRAINING KIT

You can sign up to receive a NAYZILAM® (midazolam) nasal spray, CIV training kit at [www.NAYZILAM.com](http://www.NAYZILAM.com). It includes information to help you and your loved ones practice how to give the nasal spray, so everyone can feel reassured that they know what to do during a seizure cluster emergency.

The training kit includes one training device, which is meant to model the real NAYZILAM product in look and feel, but without any actual medicine. Holding the device and pressing the plunger will simulate what it feels like to use an actual device. This training device can be reset and practiced with again and again.

*Please note: The actual device that contains medication can only be used one time.*



This information does not take the place of talking to your healthcare provider about your medical condition or treatment. If you and your family members or caregivers have any questions about NAYZILAM, ask your healthcare provider or pharmacist.



Please see additional Important Safety Information on pages 10-13 and the enclosed Patient Medication Guide in this brochure's pocket. Visit [www.NAYZILAM.com](http://www.NAYZILAM.com) to learn more about treatment with NAYZILAM.

## WHEN TO REPLACE NAYZILAM

Because seizure clusters can strike at anytime, it's always important to plan ahead. You should always have two spray devices on hand, so be sure to get a refill if you use one of the two devices that come in each box.

### REMEMBER TO...

Check the expiration date on the peel pack, back of the device, or on the bottom panel of the product box, so you know when you need to replace your nasal spray.



ucbCARES® is the hub for information on all UCB products. Reach out to our team of nurses, pharmacists, and customer service professionals to:

- Ask questions about UCB products
- Report side effects or product quality concerns
- Get clinical trial information and support
- Find information on affording your UCB medication

## CONNECT WITH US



1-844-599-CARE (2273)



[www.askucbcares.com](http://www.askucbcares.com)



## SAVINGS PROGRAM

**ELIGIBLE PATIENTS MAY PAY AS LITTLE AS \$20 FOR NAYZILAM\***

Eligible patients\* may pay as little as \$20 per box of NAYZILAM® (midazolam) nasal spray, CIV. Just present the NAYZILAM Patient Savings Card at your pharmacy when filling your prescription. Please read eligibility restrictions and terms and conditions on the back of the NAYZILAM Patient Savings Card or at [www.NAYZILAM.com/savings](http://www.NAYZILAM.com/savings).

\*Eligibility restrictions and conditions apply.

**TO CHECK YOUR ELIGIBILITY OR TO  
ACTIVATE YOUR SAVINGS CARD, VISIT:**



**www.NAYZILAM.com/savings**



\*Eligibility Criteria & Terms: This savings card is not valid for use by patients who are covered by any federal or state funded healthcare program (including, but not limited to, Medicare [Part D and Medigap] and those who are Medicare-eligible and enrolled in an employer-sponsored health plan for retirees, Medicaid, any state pharmaceutical assistance program, TRICARE, VA, or DoD), or for cash-paying patients. Offer good only in the U.S., including Puerto Rico. This card is good for use only with a valid NAYZILAM prescription consistent with the approved FDA labeling at the time the prescription is filled by the pharmacist and dispensed to the patient. The maximum annual benefit amount is \$1200 per calendar year. Void where prohibited by law, taxed, or restricted. This offer cannot be combined with any other promotional offer. UCB, Inc. reserves the right to rescind, revoke, or amend this offer without notice at any time. No cash value. Not eligible for sale, purchase, trade, or counterfeit.

**Nayzilam®** (midazolam) nasal spray

Please see additional Important Safety Information on pages 10-13 and the enclosed Patient Medication Guide in this brochure's pocket. Visit [www.NAYZILAM.com](http://www.NAYZILAM.com) to learn more about treatment with NAYZILAM.

## NOTES

This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are approximately 20 lines visible. The paper has a slight shadow on the right side, suggesting it's resting on a surface.

## This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are approximately 20 lines visible. The paper has a slight shadow on its right side, suggesting it's resting on a surface.





# STOP SEIZURE CLUSTERS QUICKLY WITH NAYZILAM

Now you can feel confident knowing that you can stop a seizure cluster when it strikes and get back to your day.

## IMPORTANT SAFETY INFORMATION

NAYZILAM is a benzodiazepine medicine. Taking benzodiazepines with opioid medicines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, breathing problems (respiratory depression), coma, and death. NAYZILAM may cause serious side effects, including: interaction with other central nervous system depressants, serious breathing problems and excessive sleepiness, suicidal thoughts or actions, impaired mental alertness including memory problems, increase in eye pressure in people with acute narrow-angle glaucoma. The most common side effects of NAYZILAM include sleepiness, nasal discomfort, headache, throat irritation and runny nose.

For more information, go to [www.NAYZILAM.com](http://www.NAYZILAM.com) or call 1-844-599-2273.

Please talk to your healthcare provider if you have any questions or concerns about NAYZILAM.

NAYZILAM® and ucbCARES® are registered trademarks of the UCB Group of Companies.

©2019 UCB, Inc., Smyrna, GA 30080. All rights reserved.

US-P-NZ-SC-1900064



Inspired by patients.  
Driven by science.