

WHEN IT COMES TO STOPPING
BACK-TO-BACK SEIZURES

NAYZILAM

READY

TAKE ON SEIZURE CLUSTERS
AND RESCUE YOUR DAY



Not an actual patient.



SEE PATIENT SAVINGS INFORMATION INSIDE

INDICATION

NAYZILAM is a prescription medicine used for the short-term treatment of seizure clusters (also known as acute repetitive seizures) that are different from a patient's usual seizure pattern in patients 12 years of age and older.

IMPORTANT SAFETY INFORMATION

NAYZILAM is a benzodiazepine medicine. Taking benzodiazepines with opioid medicines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, breathing problems (respiratory depression), coma, and death. There is a risk of abuse, misuse, and addiction with benzodiazepines, including NAYZILAM, which can lead to overdose and serious side effects including coma and death. Benzodiazepines, including NAYZILAM can cause physical dependence and withdrawal reactions, especially if you continue to use NAYZILAM daily. NAYZILAM is not intended for daily use.

Please see additional Important Safety Information and the enclosed Patient Medication Guide in this brochure's pocket. Visit NAYZILAM.com to learn more about treatment with NAYZILAM.

SEIZURE CLUSTERS CAN HAPPEN ANYTIME, OR ANYWHERE

Episodes of increased seizure activity, like having two or more seizures in a 24-hour period, are called seizure clusters or acute repetitive seizures. They are different from your usual seizure pattern. This can mean having several seizures in a row, or they can be hours apart.

The number of seizures in a cluster, type of seizure, severity, or how long they last can vary from one person to the next. Your healthcare provider can help you to identify seizure clusters, and when and how to use a rescue medication to treat them.

SEIZURE
CLUSTERS

2 SEIZURES
IN 24
HOURS
OR MORE

COMMON SEIZURE TRIGGERS



ILLNESS



STRESS



LACK OF
SLEEP



MENSTRUAL
PERIODS



MISSED
MEDICATIONS



TRAVELING



ALCOHOL

There's a lot out there that can trigger a seizure cluster—being sick, stress, not sleeping well, menstrual periods, alcohol, and even missing a medication.

Know your triggers and be prepared with a rescue therapy.



YOUR HEALTHCARE PROVIDER CAN HELP YOU IDENTIFY
SEIZURE TRIGGERS AND SHOW YOU HOW TO USE
A RESCUE NASAL SPRAY FOR SEIZURE CLUSTERS.

Rescue treatments are different from daily epilepsy medications you already take to manage your epilepsy. Keep taking your daily seizure medication as directed by your healthcare provider, and use your prescribed rescue treatment as needed to treat your seizure cluster.



WHAT IS NAYZILAM?

NAYZILAM® (midazolam) nasal spray, CIV is a prescription rescue treatment used for the short-term treatment of seizure clusters (also known as acute repetitive seizures) in patients 12 years of age and older.

A CARRY-WITH-YOU NASAL SPRAY THAT STOPS SEIZURE CLUSTERS FAST



ON-THE-GO

NAYZILAM is portable and meant to be taken with you anywhere, so you're always ready when a seizure cluster strikes.



RESULTS IN MINUTES

NAYZILAM helps stop seizure clusters within 10 minutes and can prevent additional seizures for up to 6 hours.



READY TO BE SPRAYED BY ANYONE

NAYZILAM can be given by anyone. Even if you're unable to spray NAYZILAM for yourself in the moment, another person who knows how to identify seizure clusters can spray it for you, whether or not you are conscious. No inhaling necessary!

NAYZILAM may cause serious side effects, including impaired mental alertness including memory problems. The most common side effects reported with use include:

- Sleepiness
- Headache
- Throat irritation
- Runny nose
- Nasal discomfort

These are not all the possible side effects of NAYZILAM. Call your healthcare provider for medical advice about side effects.



Please see additional Important Safety Information and the enclosed Patient Medication Guide in this brochure's pocket. Visit NAYZILAM.com to learn more about treatment with NAYZILAM.

IN A STUDY, NAYZILAM WAS SHOWN TO HELP:

- Stop seizure clusters **within minutes**
- Prevent further seizure activity for **up to 6 hours**
- Keep seizures from returning **for 24 hours**
- Get back to **feeling like yourself** within hours



I bring NAYZILAM everywhere I go. I can just grab it from my backpack or my purse and give it to myself or give it to somebody else to give to me.

— Eva, an actual patient living with seizure clusters and using NAYZILAM.



SCAN TO HEAR EVA'S STORY AT [NAYZILAM.COM](http://NAYZILAM.com)

IMPORTANT SAFETY INFORMATION

NAYZILAM may cause serious side effects, including: interaction with other central nervous system depressants, serious breathing problems and excessive sleepiness, risk of abuse, misuse, and addiction, physical dependence with withdrawal reactions, suicidal thoughts or actions, impaired mental alertness including memory problems, increase in eye pressure in people with acute narrow-angle glaucoma, and may cause your baby to have symptoms of sedation (breathing problems, sluggishness, low muscle tone), and/or withdrawal symptoms (jitteriness, irritability, restlessness, shaking, excessive crying, feeding problems). Breastfeeding during treatment with NAYZILAM may cause your baby to have sleepiness, feeding problems, and decreased weight gain. The most common side effects of NAYZILAM include sleepiness, nasal discomfort, headache, throat irritation and runny nose.



USING NAYZILAM

NAYZILAM® (midazolam) CIV can be given during or immediately after the first seizure—because you can't afford to wait to treat a seizure cluster.



ONE DOSE OF NAYZILAM = ONE SPRAY

Use one 5-mg nasal spray in one nostril. If the seizure is continuing 10 minutes after the first dose, a second dose may be used. If a second dose is used, give the second dose in the other nostril. **Do not give more than 2 doses to treat a seizure cluster.**

BE NAYZILAM READY



READY

Take NAYZILAM with you all the time. It's the **carry-with-you** rescue spray that's small enough to put in a pocket or purse.



SET

Have a personalized **Seizure Action Plan** in place so you and your friends, coworkers, and loved ones know where your NAYZILAM is stored and how and when to use it.



GO

NAYZILAM may help stop the cluster fast and **get you back to your day.**

Nayzilam®
(midazolam) nasal spray

Please see additional **Important Safety Information** and the enclosed **Patient Medication Guide** in this brochure's pocket. Visit NAYZILAM.com to learn more about treatment with NAYZILAM.

AJ is always on-the-go with friends or playing sports. He takes NAYZILAM with him everywhere he goes in case a seizure cluster occurs.

AJ, an actual patient living with seizure clusters and using NAYZILAM.



ABOUT USING NAYZILAM

It's important to know a few things before you or someone else administers NAYZILAM so everyone can feel more comfortable using it.

- NAYZILAM is not intended to be used daily. Use exactly as your healthcare provider tells you to use it (in the nose, never more than 2 doses). If you use too much of this medication, call your healthcare provider or go to the nearest emergency room right away.
- If the seizures do not stop after the nasal spray is used, get emergency medical help right away.
- Do not use this medication for more than 1 seizure cluster episode every 3 days and no more than 5 seizure cluster episodes per month.



GETTING NAYZILAM

SAVINGS PROGRAM & PATIENT SUPPORT

ELIGIBLE PATIENTS MAY PAY AS LITTLE AS \$20 FOR NAYZILAM*

Sign up for a NAYZILAM Patient Savings Card to start saving on your prescription.

HOW TO CHECK YOUR ELIGIBILITY AND GET STARTED



VISIT NAYZILAM.COM/SAVINGS



TEXT "JOINNAYZ" TO 50566

to sign up for a NAYZILAM Patient Savings Card and SMS product reminder alert[†]



CALL 1-888-786-5879



SCAN TO START SAVING AT NAYZILAM.COM

*Eligibility Criteria and Terms: This savings card is not valid for use by patients who are covered by any federally funded or state-funded healthcare program (including, but not limited to, Medicare [Part D and Medigap] and those who are Medicare-eligible and enrolled in an employer-sponsored health plan for retirees, Medicaid, any state pharmaceutical assistance program, TRICARE, VA, or DoD, or for cash-paying patients). Offer good only in the U.S., including Puerto Rico. This card is good for use only with a valid NAYZILAM prescription consistent with the approved FDA labeling at the time the prescription is filled by the pharmacist and dispensed to the patient. The maximum annual benefit amount is \$1200 per calendar year. Void where prohibited by law, taxed, or restricted. This offer cannot be combined with any other promotional offer. UCB, Inc. reserves the right to rescind, revoke, or amend this offer without notice at any time. No cash value. Not eligible for sale, purchase, trade, or counterfeit.

[†]Message and data rates may apply. Four (4) messages per month.

Nayzilam®
(midazolam) nasal spray

IMPORTANT SAFETY INFORMATION

Important Safety Information you should know about NAYZILAM® (midazolam) nasal spray, CIV

What is the most important information I should know about NAYZILAM?

NAYZILAM is a benzodiazepine medicine. Taking benzodiazepines with opioid medicines, alcohol, or other central nervous system (CNS) depressants (including street drugs) can cause severe drowsiness, breathing problems (respiratory depression), coma, and death.

Get emergency medical help right away if any of the following happens:

- shallow or slowed breathing
- breathing stops (which may lead to the heart stopping)
- excessive sleepiness (sedation)

Do not drive or operate heavy machinery until you know how taking NAYZILAM with opioids may affect you.

Risk of abuse, misuse, and addiction. There is a risk of abuse, misuse, and addiction with benzodiazepines, including NAYZILAM, which can lead to overdose and serious side effects including coma and death.

- Serious side effects including coma and death have happened in people who have abused or misused benzodiazepines, including midazolam (the active ingredient in NAYZILAM). These serious side effects may also include delirium, paranoia, suicidal thoughts or actions, seizures, and difficulty breathing. Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of these serious side effects.
- You can develop an addiction even if you use NAYZILAM as prescribed by your healthcare provider.
- Use NAYZILAM exactly as your healthcare provider prescribed.
- Do not share your NAYZILAM with other people.
- Keep NAYZILAM in a safe place and away from children.

Physical Dependence and withdrawal reactions. Benzodiazepines, including NAYZILAM can cause physical dependence and withdrawal reactions, especially if you continue to use NAYZILAM daily. NAYZILAM is not intended for daily use.

- Do not suddenly stop using NAYZILAM without talking to your healthcare provider. Stopping NAYZILAM suddenly can cause serious and life-threatening side effects, including, unusual movements, responses, or expressions, seizures, sudden and severe mental or nervous system changes, depression, seeing or hearing things that others do not see or hear, an extreme increase in activity or talking, losing touch with reality, and suicidal thoughts or actions. Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of these symptoms.
- Some people who suddenly stop benzodiazepines have symptoms that can last for several weeks to more than 12 months, including, anxiety, trouble remembering, learning, or concentrating, depression, problems sleeping, feeling like insects are crawling under your skin, weakness, shaking, muscle twitching, burning, or prickling feeling in your hands, arms, legs or feet, and ringing in your ears.
- Physical dependence is not the same as drug addiction. Your healthcare provider can tell you more about the differences between physical dependence and drug addiction.
- Do not use more NAYZILAM than prescribed or use NAYZILAM more often than prescribed.

NAYZILAM can make you sleepy or dizzy and can slow your thinking and motor skills.

- Do not drive, operate heavy machinery, or do other dangerous activities until you know how NAYZILAM affects you.
- Do not drink alcohol or take other drugs that may make you sleepy or dizzy while using NAYZILAM without first talking to your healthcare provider. When taken with alcohol or drugs that cause sleepiness or dizziness, NAYZILAM may make your sleepiness or dizziness worse.

IMPORTANT SAFETY INFORMATION (CONT'D)

Like other antiepileptic medicines, NAYZILAM may cause suicidal thoughts or actions in a very small number of people, about 1 in 500. Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- feeling agitated or restless
- acting aggressive, being angry, or violent
- attempts to commit suicide
- panic attacks
- acting on dangerous impulses
- new or worse depression
- trouble sleeping (insomnia)
- an extreme increase in activity and talking (mania)
- new or worse anxiety
- new or worse irritability
- other unusual changes in behavior or mood

How can I watch for early symptoms of suicidal thoughts or actions?

- Pay attention to any changes, especially sudden changes in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms. Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

What is NAYZILAM?

- NAYZILAM is a prescription medicine used for the short-term treatment of seizure clusters (also known as "acute repetitive seizures") in patients 12 years of age and older.
- NAYZILAM is a federally controlled substance (CIV) because it contains midazolam that can be abused or lead to dependence. Keep NAYZILAM in a safe place to prevent misuse and abuse. Selling or giving away NAYZILAM may harm others and is against the law. Tell your healthcare provider if you have abused or been dependent on alcohol, prescription drugs, or street drugs.
- It is not known if NAYZILAM is safe and effective in children under 12 years of age.

Do not use NAYZILAM if you:

- are allergic to midazolam.
- have an eye problem called acute narrow-angle glaucoma.

Before you use NAYZILAM, tell your healthcare provider about all your medical conditions, including if you:

- have a history of depression, mood problems or suicidal thoughts or behavior.
- have asthma, emphysema, bronchitis, chronic obstructive pulmonary disease, or other breathing problems.
- have kidney or liver problems.
- have congestive heart failure.
- have a history of drug or alcohol abuse
- are pregnant or plan to become pregnant. Taking NAYZILAM late in pregnancy may cause your baby to have symptoms of sedation (breathing problems, sluggishness, low muscle tone), and/or withdrawal symptoms (jitteriness, irritability, restlessness, shaking, excessive crying, feeding problems).
- are breastfeeding or plan to breastfeed. Breastfeeding during treatment with NAYZILAM may cause your baby to have sleepiness, feeding problems, and decreased weight gain.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

How should I use NAYZILAM?

- Use NAYZILAM in the nose only.
- Use NAYZILAM exactly as your healthcare provider tells you to use it and follow the Instructions for Use.
- Your healthcare provider has prescribed NAYZILAM to treat a type of seizure called a "seizure cluster".
- If the seizure cluster is continuing 10 minutes after the first dose of NAYZILAM, a second dose of NAYZILAM may be used if you have been told to do so by your healthcare provider.
- If a second dose of NAYZILAM is used, give the second dose in the other nostril.
- Do not give more than 2 doses of NAYZILAM to treat a seizure cluster.
- If the seizures do not stop after NAYZILAM is used, get emergency medical help right away.
- Do not use NAYZILAM for more than 1 seizure cluster episode every 3 days. Do not use NAYZILAM for more than 5 seizure cluster episodes per month.
- If benzodiazepines are stopped after a person takes them daily, they can cause withdrawal symptoms. Stopping benzodiazepines suddenly can cause seizures that will not stop (status epilepticus), hearing or seeing things that are not there (hallucinations), shaking, nervousness, and stomach and muscle cramps. NAYZILAM is not intended to be used daily.
- If you use too much NAYZILAM, call your healthcare provider or go to the nearest emergency room right away.

What should I avoid while using NAYZILAM?

See "What is the most important information I should know about NAYZILAM?"

What are the possible side effects of NAYZILAM?

NAYZILAM may cause serious side effects, including:

- See "What is the most important information I should know about NAYZILAM?"
- Increase in eye pressure in people with acute narrow-angle glaucoma. See "Do not use NAYZILAM if you:"

The most common side effects of NAYZILAM include:

• Sleepiness	• runny nose	• throat irritation
• headache	• nasal discomfort	


These are not all the possible side effects of NAYZILAM. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see additional patient information in the Medication Guide. This information does not take the place of talking with your healthcare provider about your condition or your treatment. For more information, go to www.NAYZILAM.com or call 1-844-599-2273.

I AM READY TO ASK FOR NAYZILAM



**SCAN TO CREATE A
DISCUSSION GUIDE
FOR YOUR NEXT APPOINTMENT**



Not an actual patient.

IMPORTANT SAFETY INFORMATION

NAYZILAM is a benzodiazepine medicine. Taking benzodiazepines with opioid medicines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, breathing problems (respiratory depression), coma, and death. There is a risk of abuse, misuse, and addiction with benzodiazepines, including NAYZILAM, which can lead to overdose and serious side effects including coma and death. Benzodiazepines, including NAYZILAM, can cause physical dependence and withdrawal reactions, especially if you continue to use NAYZILAM daily. NAYZILAM is not intended for daily use. NAYZILAM may cause serious side effects, including: interaction with other central nervous system depressants, serious breathing problems and excessive sleepiness, risk of abuse, misuse, and addiction, physical dependence with withdrawal reactions, suicidal thoughts or actions, impaired mental alertness including memory problems, increase in eye pressure in people with acute narrow-angle glaucoma. The most common side effects of NAYZILAM include sleepiness, nasal discomfort, headache, throat irritation and runny nose.

Please see the enclosed Patient Medication Guide in this brochure's pocket. For more information, go to www.NAYZILAM.com or call 1-844-599-2273. Please talk to your healthcare provider if you have any questions or concerns about NAYZILAM.

NAYZILAM® is a registered trademark of the UCB Group of Companies.
©2025 UCB, Inc., Smyrna, GA 30080. All rights reserved.
US-NZ-2500033

