

READY

TO ASK FOR ON-THE-GO RESCUE SPRAY

ARRIVE READY FOR YOUR DOCTOR APPOINTMENT

Be one step ahead of seizure clusters with NAYZILAM, the on-the-go rescue nasal spray that stops seizure clusters fast. Your doctor will want some information about your seizure patterns, triggers, and treatments before writing you a prescription.

Fill out this seizure questionnaire and be ready to discuss NAYZILAM rescue nasal spray with your doctor.



1

HAVE YOU EVER HAD 2 OR MORE SEIZURES WITHIN 24 HOURS (KNOWN AS A SEIZURE CLUSTER)?

Yes No

2

DO YOU TAKE A DAILY ANTISEIZURE MEDICATION?

Yes No

3

DO YOU HAVE A SEIZURE RESCUE MEDICATION?

Yes No

WHICH ONE(S)? _____

4

HAVE YOU NOTICED CONDITIONS THAT TRIGGER A SEIZURE CLUSTER FOR YOU? WHAT ARE THEY?

<input type="checkbox"/> Travel	<input type="checkbox"/> Lack of sleep	<input type="checkbox"/> Missed a dose	<input type="checkbox"/> None
<input type="checkbox"/> Illness	<input type="checkbox"/> Menstruation	<input type="checkbox"/> Drinking alcohol	<input type="checkbox"/> I don't know
<input type="checkbox"/> Stress	<input type="checkbox"/> Other _____		

5

HOW HAVE SEIZURES IMPACTED YOUR RELATIONSHIPS? YOUR JOB? YOUR ABILITY TO GO ABOUT EVERYDAY ACTIVITIES? YOUR STRESS LEVELS?

6

WHAT OTHER NAYZILAM TOPICS WOULD YOU LIKE TO BRING UP WITH YOUR DOCTOR?

<input type="checkbox"/> When to use NAYZILAM	<input type="checkbox"/> Using NAYZILAM with a daily antiseizure medication
<input type="checkbox"/> How to take NAYZILAM	<input type="checkbox"/> Using NAYZILAM with other medications or supplements
<input type="checkbox"/> How much NAYZILAM to take (dosing)	

DO YOU KEEP A SEIZURE DIARY?

Seizure diaries can help you track your seizure progress and identify patterns. Consider recording when you have seizures and what may have triggered them in a notebook. Be sure to take your diary with you to your doctor's appointment.

INDICATION & IMPORTANT SAFETY INFORMATION

Important Safety Information you should know about NAYZILAM® (midazolam) nasal spray, CIV

What is the most important information I should know about NAYZILAM?

NAYZILAM is a benzodiazepine medicine. Taking benzodiazepines with opioid medicines, alcohol, or other central nervous system (CNS) depressants (including street drugs) can cause severe drowsiness, breathing problems (respiratory depression), coma, and death.

Get emergency medical help right away if any of the following happens:

- shallow or slowed breathing
- breathing stops (which may lead to the heart stopping)
- excessive sleepiness (sedation)

Do not drive or operate heavy machinery until you know how taking NAYZILAM with opioids may affect you.

Risk of abuse, misuse, and addiction. There is a risk of abuse, misuse, and addiction with benzodiazepines, including NAYZILAM, which can lead to overdose and serious side effects including coma and death.

- **Serious side effects including coma and death have happened in people who have abused or misused benzodiazepines, including midazolam (the active ingredient in NAYZILAM).** These serious side effects may also include delirium, paranoia, suicidal thoughts or actions, seizures, and difficulty breathing. **Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of these serious side effects.**
- You can develop an addiction even if you use NAYZILAM as prescribed by your healthcare provider.
- Use NAYZILAM exactly as your healthcare provider prescribed.
- Do not share your NAYZILAM with other people.
- Keep NAYZILAM in a safe place and away from children.

Physical Dependence and withdrawal reactions. Benzodiazepines, including NAYZILAM can cause physical dependence and withdrawal reactions, especially if you continue to use NAYZILAM daily. NAYZILAM is not intended for daily use.

- **Do not suddenly stop using NAYZILAM without talking to your healthcare provider.** Stopping NAYZILAM suddenly can cause serious and life-threatening side effects, including, unusual movements, responses, or expressions, seizures, sudden and severe mental or nervous system changes, depression, seeing or hearing things that others do not see or hear, an extreme increase in activity or talking, losing touch with reality, and suicidal thoughts or actions. **Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of these symptoms.**
- **Some people who suddenly stop benzodiazepines have symptoms that can last for several weeks to more than 12 months,** including, anxiety, trouble remembering, learning, or concentrating, depression, problems sleeping, feeling like insects are crawling under your skin, weakness, shaking, muscle twitching, burning or prickling feeling in your hands, arms, legs or feet, and ringing in your ears.
- Physical dependence is not the same as drug addiction. Your healthcare provider can tell you more about the differences between physical dependence and drug addiction.

Do not use more NAYZILAM than prescribed or use NAYZILAM more often than prescribed.

NAYZILAM can make you sleepy or dizzy and can slow your thinking and motor skills.

- Do not drive, operate heavy machinery, or do other dangerous activities until you know how NAYZILAM affects you.
- Do not drink alcohol or take other drugs that may make you sleepy or dizzy while using NAYZILAM without first talking to your healthcare provider. When taken with alcohol or drugs that cause sleepiness or dizziness, NAYZILAM may make your sleepiness or dizziness worse.

Like other antiepileptic medicines, NAYZILAM may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.

Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

• thoughts about suicide or dying	• trouble sleeping (insomnia)
• feeling agitated or restless	• an extreme increase in activity and talking (mania)
• acting aggressive, being angry, or violent	• new or worse anxiety
• attempts to commit suicide	• new or worse irritability
• panic attacks	• other unusual changes in behavior or mood
• acting on dangerous impulses	
• new or worse depression	

How can I watch for early symptoms of suicidal thoughts or actions?

- Pay attention to any changes, especially sudden changes in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms. Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

What is NAYZILAM?

- NAYZILAM is a prescription medicine used for the short-term treatment of seizure clusters (also known as “acute repetitive seizures”) in patients 12 years of age and older.
- NAYZILAM is a federally controlled substance (CIV) because it contains midazolam that can be abused or lead to dependence. Keep NAYZILAM in a safe place to prevent misuse and abuse. Selling or giving away NAYZILAM may harm others and is against the law. Tell your healthcare provider if you have abused or been dependent on alcohol, prescription drugs, or street drugs.
- It is not known if NAYZILAM is safe and effective in children under 12 years of age.

Do not use NAYZILAM if you:

- are allergic to midazolam.
- have an eye problem called acute narrow-angle glaucoma.

INDICATION & IMPORTANT SAFETY INFORMATION (CONT'D)

Before you use NAYZILAM, tell your healthcare provider about all your medical conditions, including if you:

- have a history of depression, mood problems or suicidal thoughts or behavior.
- have asthma, emphysema, bronchitis, chronic obstructive pulmonary disease, or other breathing problems.
- have kidney or liver problems.
- have congestive heart failure.
- have a history of drug or alcohol abuse
- are pregnant or plan to become pregnant. Taking NAYZILAM late in pregnancy may cause your baby to have symptoms of sedation (breathing problems, sluggishness, low muscle tone), and/or withdrawal symptoms (jitteriness, irritability, restlessness, shaking, excessive crying, feeding problems).
- are breastfeeding or plan to breastfeed. Breastfeeding during treatment with NAYZILAM may cause your baby to have sleepiness, feeding problems, and decreased weight gain.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

How should I use NAYZILAM?

- Use NAYZILAM in the nose only.
- **Use NAYZILAM exactly as your healthcare provider tells you to use it** and follow the Instructions for Use.
- Your healthcare provider has prescribed NAYZILAM to treat a type of seizure called a "seizure cluster".
- If the seizure cluster is continuing **10 minutes** after the first dose of NAYZILAM, a second dose of NAYZILAM **may be used if you have been told to do so by your healthcare provider**.
- If a second dose of NAYZILAM is used, give the second dose in the other nostril.
- **Do not** give more than 2 doses of NAYZILAM to treat a seizure cluster.
- **If the seizures do not stop** after NAYZILAM is used, get emergency medical help right away.
- **Do not** use NAYZILAM for more than 1 seizure cluster episode every 3 days. Do not use NAYZILAM for more than 5 seizure cluster episodes per month.
- If benzodiazepines are stopped after a person takes them daily, they can cause withdrawal symptoms. Stopping benzodiazepines suddenly can cause seizures that will not stop (status epilepticus), hearing or seeing things that are not there (hallucinations), shaking, nervousness, and stomach and muscle cramps. NAYZILAM is not intended to be used daily.
- If you use too much NAYZILAM, call your healthcare provider or go to the nearest emergency room right away.

What should I avoid while using NAYZILAM?

See "What is the most important information I should know about NAYZILAM?"

What are the possible side effects of NAYZILAM?

NAYZILAM may cause serious side effects, including:

- See "What is the most important information I should know about NAYZILAM?"
- Increase in eye pressure in people with acute narrow-angle glaucoma. See "Do not use NAYZILAM if you:"

The most common side effects of NAYZILAM include:

• Sleepiness	• runny nose	• throat irritation
• headache	• nasal discomfort	

These are not all the possible side effects of NAYZILAM. Call your doctor for medical advice about side effects.

You may report side effects to FDA at 1-800-FDA-1088.

Please see additional patient information in the Medication Guide. This information does not take the place of talking with your healthcare provider about your condition or your treatment. For more information, go to www.NAYZILAM.com or call 1-844-599-2273.